

CLAIMS

1. An isolated nucleic acid fragment which
- 5 - encodes a polypeptide fragment which exhibits a substantial immunological reactivity with a rabbit polyclonal antibody raised against a polypeptide having an apparent molecular weight of 13 kDa as determined by SDS-PAGE followed by visualization, said polypeptide being derived from *Borrelia burgdorferi* B313 and being encoded by the nucleotide sequence of SEQ ID NO: 18, said rabbit polyclonal antibody exhibiting substantially no immunological reactivity with proteins from at least 95% of spirochaetes randomly selected from the group consisting of *Borrelia hermsii*, *Borrelia crocidurae*, *Borrelia anserina*, and *Borrelia hispanica*, and/or
- 10 - hybridises readily under highly stringent hybridization conditions with a DNA fragment having a nucleotide sequence selected from the group consisting of SEQ ID NO: 18, SEQ ID NO: 20, and SEQ ID NO: 22, or with a DNA fragment complementary thereto, but exhibits no substantial hybridization when the hybridization conditions are highly stringent with genomic DNA from at least 95% of spirochaetes randomly selected from the group consisting of *Borrelia hermsii*, *Borrelia crocidurae*, *Borrelia anserina*, and *Borrelia hispanica*.
- 15 -
- 20 -
2. The nucleic acid fragment according to claim 1, which encodes a polypeptide fragment comprising an amino acid sequence comprised in a polypeptide, said polypeptide
- 25 being present in whole cell preparations of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, and/or *Borrelia afzelii* ACAI but being substantially absent from whole cell preparations of at least 95% of randomly selected *Borrelia hermsii*, *Borrelia crocidurae*, *Borrelia anserina*, or *Borrelia hispanica*.
- 30 3. The nucleic acid fragment according to claim 1 or 2, which encodes a polypeptide fragment comprising at least a part of an amino acid sequence of a protein having an apparent molecular weight of 13 kDa, said protein being present in whole cell preparations of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, and/or *Borrelia afzelii* ACAI but being substantially absent from whole cell

preparations of at least 95% of randomly selected *Borrelia hermsii*, *Borrelia crocidurae*, *Borrelia anserina*, and *Borrelia hispanica*.

a 4. The nucleic acid fragment according to ~~any of the preceding claims~~ ^{claim 1}, which encodes
5 a polypeptide fragment comprising at least one epitope, said epitope being present in whole cell preparations of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI but being substantially absent from whole cell preparations of at least 95% of randomly selected *Borrelia hermsii*, *Borrelia crocidurae*, *Borrelia anserina*, and *Borrelia hispanica*.

10 5. The nucleic acid fragment according to ~~any of the preceding claims~~ ^{claim 1}, which encodes a polypeptide fragment comprising at least one epitope of a protein having an apparent molecular weight of 13 kDa, said protein being present in whole cell preparations of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia*
15 *afzelii* ACAI but being substantially absent from whole cell preparations of at least 95% of randomly selected *Borrelia hermsii*, *Borrelia crocidurae*, *Borrelia anserina*, and *Borrelia hispanica*.

20 6. The nucleic acid fragment according to ~~any of the preceding claims~~ ^{claim 1}, which encodes a polypeptide fragment which has an amino acid sequence exhibiting a sequence identity of at least 50% with SEQ ID NO: 19, SEQ ID NO: 21, or SEQ ID NO: 23, or with subsequences thereof having a length of at least 10 amino acid residues.

a 25 7. The nucleic acid fragment according to ~~any of the preceding claims~~ ^{claim 1}, wherein the nucleotide sequence has a sequence homology of at least 70% with SEQ ID NO: 18, SEQ ID NO: 20, or SEQ ID NO: 22, or with subsequences thereof having a length of at least 12 nucleotides.

a 30 8. A nucleic acid fragment according to ~~any of the preceding claims~~ ^{claim 1}, which comprises a nucleic acid fragment encoding a polypeptide fragment which comprises at least one amino acid sequence selected from the group consisting of amino acid residues 19-27, 33-36, 41-47, 95-104, 138-147 and 174-179 in SEQ ID NO: 19; amino acid residues 19-26, 32-35, 40-47, 94-101, 137-146, and 174-178 in SEQ ID NO: 21; and amino acid residues 18-26, 30-33, 39-46, 91-104, 137-145 and 173-177 in SEQ ID NO: 23.

- a
9. The nucleic acid fragment according to any of the preceding claims, which encodes a protein having an apparent molecular weight of 13 kDa which is present in whole cell preparations of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI but which is substantially absent from whole cell preparations of at least 95% of randomly selected *Borrelia hermsii*, *Borrelia crocidurae*, *Borrelia anserina*, and *Borrelia hispanica*.
10. The nucleic acid fragment according to claim 9, wherein the encoded protein is present in fraction B from *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI.
11. The nucleic acid fragment according to claim 10, wherein the encoded protein is a surface exposed protein of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI.
12. A nucleic acid fragment according to any of the preceding claims which comprises a nucleotide sequence encoding a polypeptide fragment which includes an amino acid sequence selected from the group consisting of SEQ ID NOs: 19, 21, and 23.
13. A nucleic acid fragment according to any of the preceding claims, which comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 18, 20, and 22.
14. A nucleic acid fragment according to claim 12 which consists of a nucleotide sequence encoding a polypeptide fragment consisting of an amino acid sequence selected from the group consisting of SEQ ID NOs: 19, 21, and 23.
15. A nucleic acid fragment according to claim 13, which consists of a nucleotide sequence selected from the group consisting of SEQ ID NOs: 18, 20, and 22.
16. A nucleic acid fragment according to any of claims 1-13, which encodes a fusion polypeptide.
- claim 1
- claim 1
- claim 1

09508487 31000

18. The nucleic acid fragment according to claim 16 or 17, which encodes a fusion polypeptide comprising, as a fusion partner, a polypeptide fragment which

10

- has the same amino acid sequence as at least one amino acid sequence selected from the group consisting of amino acid residues 19-27, 33-36, 41-47, 95-104, 138-147 and 174-179 in SEQ ID NO: 19; amino acid residues 19-26, 32-35, 40-47, 94-101, 137-146, and 174-178 in SEQ ID NO: 21; and amino acid residues 18-26, 30-33, 39-46, 91-104, 137-145 and 173-177 in SEQ ID NO: 23,
- is a lipoprotein selected from the outer membrane lipoprotein from *E. coli* and OspA from *Borrelia burgdorferi sensu lato*,
- is a viral protein selected from Hepatitis B surface antigen, Hepatitis B core antigen, and the influenza virus non-structural protein NS1,
- is an immunoglobulin binding protein selected from protein A, protein G, and the ZZ-peptide,
- is a T-cell epitope,
- is a B-cell epitope,
- is a bacterial fimbrial protein selected from the pilus components pilin and papA, and/or

20

25

30

- is the maltose binding protein, glutathione S-transferase, β -galactosidase, calmodulin binding protein or poly-histidine.

a 19. The nucleic acid fragment according to any of the preceding claims, which is a
5 DNA fragment.

20. A polypeptide fragment which exhibits a substantial immunological reactivity with a polyclonal rabbit antibody raised against a polypeptide having an apparent molecular weight of 13 kDa as determined by SDS PAGE followed by visualization and being
10 derived from *Borrelia burgdorferi* B313, said polypeptide comprising the amino acid sequence 1-167 of SEQ ID NO: 19, said polyclonal rabbit antibody exhibiting substantially no immunological reactivity with whole cell preparations from at least 95% of randomly selected *B. hermsii*, *B. crocidurae*, *B. anserina*, or *B. hispanica*, with the proviso that said polypeptide is essentially free from other *Borrelia*-derived antigens
15 when it is identical in amino acid sequence to a 13 kDa surface exposed polypeptide which can be extracted from *Borrelia burgdorferi sensu lato*,

the polypeptide fragment optionally being lipidated.

20 21. The polypeptide fragment according to claim 20, which comprises an amino acid sequence comprised in a polypeptide, said polypeptide being present in whole cell preparations of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI but being substantially absent from whole cell preparations of at least 95% of randomly selected *Borrelia hermsii*, *Borrelia crocidurae*,
25 *Borrelia anserina*, or *Borrelia hispanica*.

a 22. The polypeptide fragment according claim 20 or 21, which comprises at least a part of the amino acid sequence of a protein having an apparent molecular weight of 13 kDa, said protein being present in whole cell preparations of *Borrelia burgdorferi*
30 B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI but being substantially absent from whole cell preparations of at least 95% of randomly selected *Borrelia hermsii*, *Borrelia crocidurae*, *Borrelia anserina*, and *Borrelia hispanica*.

claim 20

23. A polypeptide fragment according to ~~any of claims 20-22~~, which comprises at least one epitope, said epitope being present in whole cell preparations of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI but being substantially absent from whole cell preparations of at least 95% of randomly selected *Borrelia hermsii*, *Borrelia crocidurae*, *Borrelia anserina*, and *Borrelia hispanica*.

claim 20

24. A polypeptide fragment according to ~~any of claims 20-23~~, which comprises at least one epitope of a protein having an apparent molecular weight of 13 kDa, said protein being present in whole cell preparations of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI but being substantially absent from whole cell preparations of at least 95% of randomly selected *Borrelia hermsii*, *Borrelia crocidurae*, *Borrelia anserina*, and *Borrelia hispanica*.

claim 20

25. The polypeptide fragment according to ~~claim any of claims 20-24~~, which comprises at least one amino acid sequence selected from the group consisting of amino acid residues 19-27, 33-36, 41-47, 95-104, 138-147 and 174-179 in SEQ ID NO: 19; amino acid residues 19-26, 32-35, 40-47, 94-101, 137-146, and 174-178 in SEQ ID NO: 21; and amino acid residues 18-26, 30-33, 39-46, 91-104, 137-145 and 173-177 in SEQ ID NO: 23.

claim 20

26. The polypeptide fragment according to ~~any of claims 20-25~~, which has an amino acid sequence identical to that of a protein having an apparent molecular weight of 13 kDa and being present in whole cell preparations of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI.

27. The polypeptide fragment according to ~~claim 26~~, wherein the protein is present in fraction B from *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI.

28. The polypeptide fragment according to ~~claim 26 or 27~~, wherein the protein is a surface exposed protein of *Borrelia burgdorferi* B31, *Borrelia burgdorferi* B313, *Borrelia garinii* IP90, or *Borrelia afzelii* ACAI.

claim 20
~~claim 20-2~~

10

15

33. A fusion polypeptide comprising as a first fusion partner the polypeptide according to any of ~~claims 20-32~~ ^{claim 20}.

25

30

which has the same amino acid sequence as at least one amino acid sequence selected from the group consisting of amino acid residues 19-27, 33-36, 41-47, 95-104, 138-147 and 174-179 in SEQ ID NO: 19; amino acid residues 19-26, 32-35, 40-47, 94-101, 137-146, and 174-178 in SEQ ID NO: 21; and amino acid residues 18-26, 30-33, 39-46, 91-104, 137-145 and 173-177 in SEQ ID NO: 23.

- Sub
Q2

SECRET

Sub
94

2

Sub
a3

- 5 41. A vector according to claim 40, wherein the promoter drives expression in a eukaryotic cell.
42. A vector according to claim 40 or 41, wherein the leader peptide enables secretion from or integration into the membrane of a mammalian cell.
- 10 43. A transformed cell carrying the vector of any of claims 36-42 and capable of replicating the nucleic acid fragment according to any of claims 1-19.
44. A transformed cell according to claim 43, which is a microorganism selected from a bacterium, a yeast, a protozoan, or a cell derived from a multicellular organism
- 15 selected from a fungus, an insect cell, a plant cell, and a mammalian cell.
45. A transformed cell according to claim 44 which is a bacterium of the genus *Escherichia*, *Bacillus* or *Salmonella*.
- 20 46. A transformed cell according to claim 45, which is an *E. coli* cell.
47. A stable cell line producing the polypeptide according to any of claims 20-35, which carries the vector according to any of claims 36-42 and which expresses the
- 25 nucleic acid fragment according to any of claims 1-19.
48. A method of preparing a polypeptide fragment as defined in any of claims 1-19, the method comprising
- 30 - culturing the transformed cell according to any of claims 43-46 or the stable cell line according to claim 47 under conditions facilitating the expression of the polypeptide fragment thereby, and

- or

5

49. A method according to claim 48, wherein the post-translational modifications
10 involve lipidation, glycosylation, cleavage and/or elongation.

đám 20

50. A vaccine comprising an amount of the polypeptide fragment according to ~~any of~~
~~claims 20-35 or of the polypeptide fragment prepared by the method according to~~
~~claim 48 or 49~~, the amount of the polypeptide fragment being effective to confer sub-
15 stantially increased resistance to infections with *Borrelia burgdorferi sensu lato* in an
animal, including a human being, the polypeptide fragment being formulated in com-
bination with a pharmaceutically acceptable carrier, diluent or vehicle, and the vaccine
optionally further comprising an adjuvant.

51. A vaccine according to claim 50, wherein the pharmaceutically acceptable carrier, vehicle, or diluent is selected from the group consisting of sterile water, physiological saline, glucose, polyalkylene glycols, and triglycerides; and wherein the adjuvant is selected from the group consisting of aluminium hydroxide or phosphate (alum), synthetic polymers of sugars (Carbopol), bacterial cells such as *C. parvum* or endotoxins or lipopolysaccharide components of gram-negative bacteria, physiologically acceptable oil vehicles such as mannide mono-oleate (Aracel A), a perfluorocarbon (Fluosol-DA).

52. A vaccine according to claim 50 or 51, wherein the amount of the polypeptide
30 fragment is in the range of 1-1000 μg per dose unit, such as between 2 and 750 μg ,
between 5 and 500 μg , between 7.5 and 250 μg , between 10 and 150 μg , between
10 and 100 μg , between 10 and 75 μg , and between 10 and 50 μg .

Sub
a6

53. A live vaccine comprising a non-pathogenic microorganism carrying and being capable of expressing the nucleic acid fragment according to any of claims 1-19 so as to produce the polypeptide fragment according to any of claims 20-35, the live vaccine being effective in conferring increased resistance to infection with *Borrelia burgdorferi sensu lato* in an animal, including a human being.

54. The live vaccine according to claim 53, wherein the non-pathogenic microorganism is selected from the group consisting of *Mycobacterium bovis* BCG, *Salmonella typhi*, *Salmonella typhimurium*, *Salmonella paratyphi*, *Staphylococcus aureus*, and *Listeria monocytogenes*.

55. A combination vaccine comprising

an amount of the polypeptide fragment according to ~~any of claims 20-35 or of~~ ^{claim 20} ~~the polypeptide fragment prepared by the method according to claim 48 or 49,~~

the amount of the polypeptide fragment being effective to confer substantially increased resistance to infections with *Borrelia burgdorferi sensu lato* in an animal, including a human being;

and

at least one further *Borrelia* antigen,

the polypeptide fragment and the antigen being formulated in combination with a pharmaceutically acceptable carrier, vehicle, or diluent and the vaccine optionally further comprising an adjuvant.

56. A combination vaccine according to claim 55, wherein the at least one further *Borrelia* antigen is selected from the group consisting of OspA, OspB, OspC, OspD, OspE, OspF, OspG, PC, Oms28, Oms45, Oms 66, decorin binding protein (dbp), LpLA7, EppA, T5, S1, 26 kDa, 39 kDa, 66 kDa, 79 kDa, 85 kDa, and 110 kDa antigen.

57. A combination vaccine comprising at least two non-identical polypeptide fragments according to ~~any of claims 20-35~~ ^{claim 20} or ~~at least two non-identical polypeptide~~

~~fragments prepared by the method according to claim 48 or 49~~, the vaccine comprising an amount of the polypeptide fragments effective to confer substantially

- 5 increased resistance to infections with *Borrelia burgdorferi sensu lato* in an animal, including a human being, in combination with a pharmaceutically acceptable carrier, vehicle, or diluent, the vaccine optionally further comprising an adjuvant.

58. A vaccine comprising the nucleic acid fragment according to any of ~~claims 1-19~~ or

- 10 a vector according to any of claims 36-42, the vaccine effecting *in vivo* expression of antigens by an animal, including a human being, to whom the vaccine has been administered, the amount of expressed antigens being effective to confer substantially increased resistance to infections with *Borrelia burgdorferi sensu lato* in an animal, including a human being.

15

59. A diagnostic composition adapted for the determination of *Borrelia burgdorferi sensu lato* in a sample, the composition comprising the polypeptide fragment

according to ~~any of claims 20-35~~ ^{claim 20} or ~~the polypeptide fragment prepared by the method according to claim 48 or 49~~, the amount of the polypeptide fragment being effective

- 20 to detectably react with antibodies present in the sample, the antibodies being directed against *Borrelia burgdorferi sensu lato*, the composition optionally comprising a detectable label.

60. A diagnostic composition adapted for the determination of *Borrelia burgdorferi*

- 25 *sensu lato* in a sample, the composition comprising an amount of the nucleic acid fragment according to ~~any of claims 1-19~~ ^{claim 1} which is effective to detectably bind to a nucleic acid fragment from *Borrelia burgdorferi sensu lato* present in the sample, the composition optionally comprising a detectable label.

- 30 61. A method of immunizing an animal, including a human being, against infections with *Borrelia burgdorferi sensu lato*, the method comprising administering to the animal an immunogenically effective amount of the vaccine according to ~~any of claims~~ ^{claim 50}

~~50-58.~~

5 bound antibody resulting from the administration or incubation.

10 acids resulting from the incubation.

15 presence of amplified nucleic acid which is specific for *Borrelia burgdorferi sensu lato*.

65. A diagnostic kit comprising

- 20

30

66. A method for the preparation of an immunological composition such as a vaccine, the method comprising the steps of admixing

SECRET

Sub
as

claim 20

- a
- a polypeptide fragment according to ~~any of claims 20-35 or prepared according to the method of claim 48 or 49,~~
 - a pharmaceutically acceptable carrier, vehicle, or diluent, and optionally
 - 5 - an adjuvant.

Add
a9add
1
O

000700 28180560 09508487 031000